



# Soils Map



State: **Minnesota**  
 County: **Kandiyohi**  
 Location: **32-119N-35W**  
 Township: **Willmar**  
 Acres: **80.47**  
 Date: **10/2/2018**



Maps Provided By:  
  
 CUSTOMIZED ONLINE MAPPING  
 © Agridata, Inc. 2018 www.AgridataInc.com



Soils data provided by USDA and NRCS.

Area Symbol: MN067, Soil Area Version: 16

Code	Soil Description	Acres	Percent of field	PI Legend	Non-Irr Class *c	Productivity Index	Bromegrass alfalfa	Bromegrass alfalfa hay	Cor n	Oats	Soybeans	Spring wheat	NCCPI Soybeans
927	Harps-Glencoe-Seaforth complex, 0 to 3 percent slopes	53.58	66.6%		llw	90							79
L163A	Okoboji silty clay loam, 0 to 1 percent slopes	9.83	12.2%		lllw	86							77
1900	Okoboji-Canisteo depressional complex, 0 to 1 percent slopes	5.75	7.1%		lllw	86							78
423	Seaforth loam, 1 to 3 percent slopes	4.17	5.2%		lls	95							84
897B	Seaforth-Swanlake loams, 2 to 6 percent slopes	2.85	3.5%		lls	90	5.4	4.2	165	85	50	52	67
447	Harpster silty clay loam	2.12	2.6%		llw	90	5.1	3.8	165	82	50	49	78
954C	Swanlake-Ves loams, 6 to 12 percent slopes	1.33	1.7%		llle	78	5.1	3.9	142	75	43	45	57
L201A	Normania loam, 1 to 3 percent slopes	0.59	0.7%		le	99							82
999B	Ves-Swanlake-Hawick complex, 2 to 6 percent slopes	0.25	0.3%		lle	74	5.1	3.9	135	80	41	48	49
<b>Weighted Average</b>						<b>89.3</b>	<b>0.4</b>	<b>0.3</b>	<b>13</b>	<b>6.7</b>	<b>3.9</b>	<b>4</b>	<b>78.1</b>

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.